



**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

## CABRIO® FUNGICIDE

**ACTIVE CONSTITUENT: 250 g/L PYRACLOSTROBIN**  
**SOLVENT: 705 g/L HYDROCARBONS, LIQUID**

GROUP	<b>11</b>	FUNGICIDE
-------	-----------	-----------

For the control of leaf speckle and leaf spot in bananas and downy and powdery mildew in grapevines, husk spot in macadamia and rust in almond as specified in the Directions for Use Table.

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE.**

CONTENTS: 5L

BASF Australia Ltd ABN 62 008 437 867  
Level 12, 28 Freshwater Place Southbank VICTORIA 3006

® Registered trademark of BASF

APVMA Approval No.: 55733/58739

**STORAGE AND DISPOSAL**

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

**SAFETY DIRECTIONS**

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) a washable hat elbow-length PVC gloves, and face shield or goggles. When using a pre-prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) a washable hat and elbow-length PVC gloves. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If swallowed, do NOT induce vomiting. Give a glass of water.

**SAFETY DATA SHEET**

Additional information is listed in the Safety Data Sheet.

**CONDITIONS OF SALE**

All conditions and warranties rights and remedies implied by law or arising in contract or tort whether due to the negligence of BASF Australia Ltd or otherwise are hereby expressly excluded so far as the same may legally be done provided however that any rights of the Buyer pursuant to non-excludable conditions or warranties of the Trade Practices Act 1974 or any relevant legislation of any State are expressly preserved but the liability of BASF Australia Ltd or any intermediate Seller pursuant thereto shall be limited if so permitted by the said legislation to the replacement of the goods sold or the supply of equivalent goods and all liability for indirect or consequential loss or damage of whatsoever nature is expressly excluded. This product must be used or applied strictly in accordance with the instructions appearing hereon. This product is solely sold for use in Australia and must not be exported without the prior written consent of BASF Australia Ltd.

APVMA Approval No: 55733/58739

**CABRIO FUNGICIDE****250 g/L PYRACLOSTROBIN****HAZARD STATEMENT:**

Toxic if swallowed. May be fatal if swallowed and enters airways. Harmful if inhaled. May cause respiratory irritation. Suspected of causing cancer. Very toxic to aquatic life with long lasting effects.

**PRECAUTIONARY STATEMENTS (PREVENTION):**

Obtain specialist advice before use. Do not handle until all safety precautions have been read and understood. Avoid breathing mist. Use only outdoors or in a wellventilated area.

IF INHALED: Remove person to fresh air and keep comfortable for breathing. IF IN EYES: Rinse continuously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If exposed or concerned: Get medical advice/attention. Rinse mouth. If skin irritation occurs: Get medical advice/attention. If eye irritation persists: Call a POISON CENTRE or doctor/physician.

**PRECAUTIONARY STATEMENTS (STORAGE):**

Store locked up.

Batch No:

Date of Manufacture:

BASF Australia Ltd  
ABN 62 008 437 867  
Level 12, 28 Freshwater Place  
Southbank VICTORIA 3006

FOR SPECIALIST ADVICE IN AN EMERGENCY ONLY PHONE 1800 803 440 TOLL FREE-ALL HOURS-AUSTRALIA WIDE

Disclosure log

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# CABRIO® FUNGICIDE

**ACTIVE CONSTITUENT: 250 g/L PYRACLOSTROBIN**  
**SOLVENT: 705 g/L HYDROCARBONS, LIQUID**

GROUP	<b>11</b>	FUNGICIDE
-------	-----------	-----------

For the control of leaf speckle and leaf spot in bananas and downy and powdery mildew in grapevines, husk spot in macadamia and rust in almond as specified in the Directions for Use Table.

THIS LEAFLET IS PART OF THE LABEL.

CONTENTS: 5L

BASF Australia Ltd ABN 62 008 437 867  
Level 12, 28 Freshwater Place Southbank VICTORIA 3006

® Registered trademark of BASF

APVMA Approval No.: 55733/58739

## DIRECTIONS FOR USE

### RESTRAINTS

- DO NOT use by aerial application in grapevines, almond and macadamia crops.  
 DO NOT apply within 48 hours of heavy rainfall.  
 DO NOT irrigate to the point of run-off.  
 DO NOT apply to water-logged soil.

### MANDATORY NO-SPRAY ZONES

DO NOT apply if there are aquatic and wetland areas including aquacultural ponds, surface streams and rivers downwind from the application area and within the **mandatory no-spray zones** shown in Table 1 below:

Table 1 – No-Spray Zones for Protection of the Aquatic	
FOR AERIAL APPLICATION	
Crop	Downwind Mandatory No-
Bananas	30 metres
FOR GROUND APPLICATION	
Crop	Downwind Mandatory No-
Bananas	10 metres
Grapes	10 metres
Macadamias	120 metres
Almonds	160 metres

CROP	DISEASE	RATE	WHP (Days)	CRITICAL COMMENTS
Bananas (bagged)	Leaf speckle ( <i>Mycosphaerella musae</i> ), leaf spot ( <i>Mycosphaerella musicola</i> ) Black Sigatoka ( <i>Mycosphaerella fijiensis</i> )	300 to 400 mL plus 3 to 5 L of water miscible oil/ha.	-	Maintain good crop hygiene by regularly removing old and diseased leaves to reduce the source of disease inoculum. DO NOT use on bananas unless bunch covers are in place. Ground application and aerial application: Apply by airblast sprayer or misting machine. Use the higher rates under conditions favouring disease infection. Apply up to four sprays per season, as part of a preventive disease control programme, commencing early in the season and using alternative mode-of-action products in between CABRIO applications. Do NOT apply consecutive sprays of CABRIO or other Group 11 fungicides. This use is subject to an CropLife Australia resistance management strategy.

CROP	DISEASE	RATE	WHP (Days)	CRITICAL COMMENTS
Grapevines	Downy mildew ( <i>Plasmopara viticola</i> ), powdery mildew ( <i>Uncinula necator</i> ),	Dilute spray 40 mL/100 L water  Concentrate spray Refer to the "Application" section.	21	Also see 'CAUTION' section re export commodities. Apply up to three sprays per season as part of a complete disease control programme. Ideally, apply in a block of three sprays at 10 to 14 day intervals, commencing at flowering. Use the shorter intervals under conditions favouring disease infection. Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. This use is subject to an CropLife Australia resistance management strategy.
Macadamia	Husk spot ( <i>Pseudocercospora macadamiae</i> )	40mL/100L in maximum spray volume of 2000L/ha	-	Apply up to two sprays per season as part of a complete disease control program. Commence CABRIO application at match head timing stage and repeat application at 14 to 28 days later. Ensure that fungicides from an alternative chemical group are included in the spray program each season.
Almond	Rust ( <i>Tranzschelia discolor</i> )	40mL/100L	-	Apply up to two sprays per season as part of a complete disease control program. Commence application at flowering and repeat application at 10 to 14 days later. Ensure that fungicides from an alternative chemical group are included in the spray program each season.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

#### **WITHHOLDING PERIODS:**

#### **HARVEST:**

ALMOND, BANANAS (BAGGED), MACADAMIA - NOT REQUIRED WHEN USED AS DIRECTED.

GRAPES - DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION

#### **GENERAL INSTRUCTIONS**

##### **MIXING**

To ensure even mixing, half-fill the spray tank with clean water and add the required amount of product. If required, add compatible products and agitate thoroughly, then add the remainder of the water. Agitate again before spraying commences.

##### **APPLICATION**

##### **Bananas:**

Ground application: Apply by airblast sprayer or misting machine to ensure even coverage.

Aerial application: Apply in a minimum volume of 20 L/ha.

**Grapevines/ Macadamia/ Almond:****Dilute Spraying**

- ◆ Use a sprayer designed to apply high spray volumes, up to the point of run-off and matched to the crop being sprayed.
- ◆ Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient spray solution to cover the crop to the point of run-off. Avoid excessive run-off.
- ◆ The required spray volume to achieve point of run off may be determined by applying different test volumes, using different settings on the sprayer, or from industry guidelines or other expert advice.
- ◆ Add the amount of product specified in the Directions for Use Table for each 100 L of water. Spray to the point of run-off.
- ◆ The required dilute spray volume to achieve point of run off will change and the sprayer set up and operation may also need to be changed, as the crop grows.

**Concentrate Spraying**

- ◆ Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies spray volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- ◆ Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen spray volume.
- ◆ Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- ◆ The mixing rate for concentrate spraying can then be calculated in the following way:

**EXAMPLE ONLY**

1. Dilute spray volume as determined above: For example 1500 L/ha
  2. Your chosen concentrate spray volume: For example 500 L/ha
  3. The concentration factor in this example is: 3 X (i.e.  $1500 \text{ L} \div 500 \text{ L} = 3$ )
  4. If the dilute label rate is 20 mL/100 L, then the concentrate rate becomes 3 x 20 (that is 60 mL of product per 100 L water for concentrate spraying).
- ◆ The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows.
  - ◆ For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

**COMPATIBILITY**

For information on compatibility please contact your local BASF representative.

**FUNGICIDE RESISTANCE WARNING****GROUP 11 FUNGICIDE**

CABRIO Fungicide is a member of the strobilurin group of fungicides. For fungicide resistance management, CABRIO Fungicide is a Group 11 fungicide. Some naturally-occurring individual fungi resistant to CABRIO Fungicide and other Group 11 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by CABRIO Fungicide or other Group 11 fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, BASF Australia Ltd accepts no liability for any losses that may result from the failure of CABRIO Fungicide to control resistant fungi.

**PRECAUTIONS**

DO NOT use human flaggers/markers unless they are protected by engineering controls such as enclosed cabs.

**RE-ENTRY PERIOD**

Do NOT allow entry into treated areas until the spray has dried. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

**CAUTION****EXPORT OF TREATED FRUIT OR WINE**

Growers should note that Maximum Residue Limits (MRLs) or import tolerances do not exist in all markets for fruit treated with CABRIO Fungicide. Additionally, some export markets have established MRLs different to those in Australia. If you are growing fruit for export (either fresh, dried or for wine production), please check with BASF Australia Ltd or the Australian Wine Research Institute <http://www.awri.com.au> for the latest information on MRLs and import tolerances BEFORE using CABRIO Fungicide.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

HIGHLY TOXIC TO AQUATIC ORGANISMS. Do NOT contaminate streams, rivers or waterways with the chemical or used containers.

**STORAGE AND DISPOSAL**

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

**SAFETY DIRECTIONS**

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) a washable hat elbow-length PVC gloves, and face shield or goggles. When using a pre-prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) a washable hat and elbow-length PVC gloves. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length PVC gloves and face shield or goggles. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat and elbow-length PVC gloves. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If swallowed, do NOT induce vomiting. Give a glass of water.

**SAFETY DATA SHEET**

Additional information is listed in the Safety Data Sheet.

**CONDITIONS OF SALE**

All conditions and warranties rights and remedies implied by law or arising in contract or tort whether due to the negligence of BASF Australia Ltd or otherwise are hereby expressly excluded so far as the same may legally be done provided however that any rights of the Buyer pursuant to non-excludable conditions or warranties of the Trade Practices Act 1974 or any relevant legislation of any State are expressly preserved but the liability of BASF Australia Ltd or any intermediate Seller pursuant thereto shall be limited if so permitted by the said legislation to the replacement of the goods sold or the supply of equivalent goods and all liability for indirect or consequential loss or damage of whatsoever nature is expressly excluded. This product must be used or applied strictly in accordance with the instructions appearing hereon. This product is solely sold for use in Australia and must not be exported without the prior written consent of BASF Australia Ltd.

APVMA Approval No: 55733/58739

**CABRIO FUNGICIDE****250 g/L PYRACLOSTROBIN****HAZARD STATEMENT:**

Toxic if swallowed. May be fatal if swallowed and enters airways. Harmful if inhaled. May cause respiratory irritation. Suspected of causing cancer. Very toxic to aquatic life with long lasting effects.



**PRECAUTIONARY STATEMENTS (PREVENTION):**

Obtain specialist advice before use. Do not handle until all safety precautions have been read and understood. Avoid breathing mist. Use only outdoors or in a wellventilated area.

IF INHALED: Remove person to fresh air and keep comfortable for breathing. IF IN EYES: Rinse continuously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If exposed or concerned: Get medical advice/attention. Rinse mouth. If skin irritation occurs: Get medical advice/attention. If eye irritation persists: Call a POISON CENTRE or doctor/physician.

**PRECAUTIONARY STATEMENTS (STORAGE):**

Store locked up.

Batch No:

Date of Manufacture:

® = Registered trademark of BASF



The Chemical Company

BASF Australia Ltd  
ABN 62 008 437 867  
Level 12, 28 Freshwater Place  
Southbank VICTORIA 3006

FOR SPECIALIST ADVICE IN AN EMERGENCY ONLY PHONE 1800 803 440 TOLL FREE-ALL HOURS-AUSTRALIA WIDE.

Disclosure log

**POISON**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# Bulldock® 25 EC

**INSECTICIDE**

**ACTIVE CONSTITUENT: 25 g/L BETA-CYFLUTHRIN**  
**SOLVENT: 803 g/L LIQUID HYDROCARBONS**

**GROUP 3A INSECTICIDE**

For control of certain insect pests of avocados and macadamias as specified in the **DIRECTIONS FOR USE** table

**DIRECTIONS FOR USE**

**RESTRAINT**

**DO NOT** apply if rain is expected within 6 hours.

CROP	PEST	RATE	WHP	CRITICAL COMMENTS
Avocados	Fruitspotting bug	<b>Dilute spraying</b> 25 to 50 mL/100 L water  <b>Concentrate spraying</b> Refer to the Application section	7 days	Apply by dilute or concentrate spraying equipment. Regularly monitor avocado trees for fruitspotting bug presence and spray when numbers exceed action thresholds. Use the higher rate for high levels of infestation or likelihood of continuous reinfestation. Repeat application when numbers exceed the action threshold. Apply a maximum of 4 sprays with a minimum of 21 days between consecutive sprays. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. Can be used at up to 5 X concentrate. Do not use at concentrate rates greater than 250 mL/100 L of water (i.e. 5X concentrate).
Macadamia (Qld, NSW, ACT, WA, only)	Macadamia nut borer	<b>Dilute spraying</b> 50 mL/100 L  <b>Concentrate spraying</b> Refer to the Application section	7 days	Apply by dilute or concentrate spraying equipment. Spray to thoroughly cover nuts when pest numbers indicate, or at 2 to 3 weekly intervals during the period when pests are normally active. Use the shorter interval and the higher rate during wet weather or severe infestations. Avoid season-long application of the same insecticide. Good pest management involves the use of alternative chemical groups to avoid resistance build-up.
	Fruitspotting bug	<b>Dilute spraying</b> 25 or 50 mL/100 L  <b>Concentrate spraying</b> Refer to the Application section		Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. Can be used at up to 5 X concentrate. Do not use at concentrate rates greater than 250 mL/100 L of water (i.e. 5X concentrate).

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIODS (WHP)**

**Avocados, Macadamias: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION**

## CONTENT

DIRECTIONS FOR USE.....	2
RESTRAINT.....	2
WITHHOLDING PERIODS (WHP).....	2
EXPORT OF TREATED PRODUCE.....	3
GENERAL INSTRUCTIONS.....	3
Mixing.....	3
Application.....	3
Special instructions for Avocados and Macadamias.....	3
COMPATIBILITY.....	3
INSECTICIDE RESISTANCE WARNING.....	4
PROTECTION OF LIVESTOCK.....	4
PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT.....	4
STORAGE AND DISPOSAL.....	4
SAFETY DIRECTIONS.....	4
FIRST AID.....	4
SAFETY DATA SHEET.....	4
EXCLUSION OF LIABILITY.....	4
GHS STATEMENTS.....	5

Disclosure log

## EXPORT OF TREATED PRODUCE

Growers should note that suitable MRLs or import tolerances may not be established in all markets for produce treated with Bulldock. If you are growing for export, please check with Bayer Crop Science for the latest information on MRLs and import tolerances before using Bulldock.

## GENERAL INSTRUCTIONS

### Mixing

Add the required quantity of Bulldock 25 EC to water in the spray tank and mix thoroughly.

### Application

Bulldock 25 EC can be applied by air or ground equipment. In common with other non-systemic insecticides adequate coverage with Bulldock 25 EC is essential for maximum effectiveness.

When applying by air use a minimum of 20 litres of spray mixture per hectare. Spray in crosswinds. Do not spray in calms or when wind is light and variable in direction unless smoke indicators show spray is entering the crop uniformly. Spray application should be carried out during the cooler parts of the day or night.

A spraydrift minimisation strategy should be employed at all times when aerially applying sprays to, or near, sensitive areas. The strategy envisaged is exemplified by the cotton industry's Best Management Practice Manual.

## Special instructions for Avocados and Macadamias

### Dilute Spraying

- ◆ Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.
- ◆ Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off.
- ◆ The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- ◆ Add the amount of product specified in the Directions for Use table for each 100 L of water. Spray to the point of run-off.
- ◆ The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

### Concentrate Spraying

- ◆ Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- ◆ Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- ◆ Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- ◆ The mixing rate for concentrate spraying can then be calculated in the following way:

#### **EXAMPLE ONLY**

1. Dilute spray volume as determined above: For example 1500 L/ha
  2. Your chosen concentrate spray volume: For example 500 L/ha
  3. The concentration factor in this example is: 3X (i.e  $1500 \text{ L} \div 500 \text{ L} = 3$ )
  4. If the dilute label rate is 25 mL/100 L, then the concentrate rate becomes  $3 \times 25$ , that is 75 mL/100 L of concentrate spray.
- ◆ The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows.
  - ◆ Do not use a concentrate rate higher than that specified in the Critical Comments.
  - ◆ For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

## COMPATIBILITY

Bulldock 25 EC is compatible with most commonly used insecticides and fungicides. DO NOT mix concentrates together but add each to the spray tank separately. As formulations of other manufacturers' products are beyond the control of Bayer CropScience Pty Ltd, all mixtures should be tested prior to mixing commercial quantities. As changes in climatic conditions can alter the sensitivity of plants to mixtures of sprays, Bayer Crop Science cannot be responsible for the behaviour of such mixtures.

## **INSECTICIDE RESISTANCE WARNING**

### **GROUP 3A INSECTICIDE**

Some naturally occurring insect biotypes resistant to Bulldock 25 EC and other Group 3A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Bulldock 25 EC or other Group 3A insecticides are used repeatedly. The effectiveness of Bulldock 25 EC on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Bayer CropScience Pty Ltd accepts no liability for any losses that may result from the failure of Bulldock 25 EC to control resistant insects. Bulldock 25 EC may be subject to specific resistance management strategies. For further information contact your local supplier, Bayer Crop Science representative or local agricultural department agronomist. Application of Bulldock 25 EC to synthetic pyrethroid resistant larvae longer than 5 mm will not only be ineffective but it may increase the level of synthetic pyrethroid resistance.

### **PROTECTION OF LIVESTOCK**

Dangerous to bees. DO NOT spray any plants in flower while bees are foraging.

### **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

### **STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. Do not reuse empty container for any other purpose.

### **SAFETY DIRECTIONS**

Poisonous if swallowed. Avoid inhaling vapour or spray mist. Will damage eyes and skin. Avoid contact with eyes and skin. When opening container and preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and washable hat, elbow length PVC gloves and face shield or goggles. If clothing becomes contaminated with product remove clothing immediately. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

### **FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre (telephone 13 11 26). If swallowed do NOT induce vomiting. Give a glass of water. Remove from contaminated area. Apply artificial respiration if not breathing.

### **SAFETY DATA SHEET**

Additional information is listed in the Safety Data Sheet, which can be obtained from [www.crop.bayer.com.au](http://www.crop.bayer.com.au).

### **EXCLUSION OF LIABILITY**

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

Bulldock® is a Registered Trademark of the Bayer Group.

APVMA Approval No.: 40422/61825a

FOR 24 HOUR SPECIALIST ADVICE  
IN EMERGENCY ONLY  
PHONE 1800 033 111

## GHS STATEMENTS

**•Harmful if inhaled. •Causes skin irritation and serious eye damage. •Suspected of causing cancer.**  
**•May be fatal if swallowed and enters airways.**  
•Do not handle until all safety precautions have been read and understood. •IF SWALLOWED: Immediately call a POISON CENTRE or doctor/ physician. Do NOT induce vomiting. •IF ON SKIN: Wash with plenty of soap and water. If skin irritation occurs: Get medical advice/attention. •IF INHALED: Remove person to fresh air and keep comfortable for breathing. Call a POISON CENTRE or doctor/physician if you feel unwell. •IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTRE or doctor/physician. •Store locked up.

Disclosure log











Agrofarm 100 G





Disclosure Log